

Page 1 of 4

**FORM PTO-1449 U.S. Department of Commerce  
Patent and Trademark Office**

Attorney Docket Number  
3477.91

Serial No.  
09/744,167

**RECEIVED**

**SEP 30 2002**

**LIST OF DOCUMENTS CITED BY APPLICANT**

(Use several sheets if necessary)

Applicants:

Jeffrey L. Wrana

**TECH CENTER 1600/2900**

Filing Date  
January 19, 2001

Group

**U. S. PATENT DOCUMENTS**

Examiner Initial	Document Number	Date	Name	Class	Subclass	Filing Date if Appropriate

**FOREIGN PATENT DOCUMENTS**

Document Number	Date	Country	Class	Subclass	Translation Yes   No

**OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)**

MW	1.	Abdollah, S., Macias-Silva, M., Tsukazaki, T., Hayashi, H., Attisano, L., and Wrana, J.L. (1997). TBRI phosphorylation of Smad2 on Ser 465 and 467 is required for Smad2/Smad4 complex formation and signaling. J. Biol. Chem. 272, 27678-27685.
.	2.	Attisano, L. and Wrana, J.L. (1998). Mads and Smads in TGFB signaling. Curr. Op. Cell Biol. 10, 188-194.
.	3.	B. Meckelein et al., (1998), Mol. Brain Research, v. 55, pp. 181-197.
MW	4.	Burd et al., (1998), Mol. Cell., 2, 157-162.

EXAMINER  
\*EXAMINER

DATE CONSIDERED

Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449 U.S. Department of Commerce  
Patent and Trademark Office

Attorney Docket Number  
3477.91

Serial No.  
09/744,167

LIST OF DOCUMENTS CITED BY APPLICANT

(Use several sheets if necessary)



RECEIVED

SEP 30 2002

Applicants:

Jeffrey L. Wrana

TECH CENTER 1600/2900

Filing Date  
January 19, 2001

Group

123	5.	Chen, Y., Bhushan, A., and Vale, W. (1997b). Smad8 mediates the signaling of the receptor serine kinase. Proc. Natl. Acad. Sci. USA 94, 12938-12943.
	6.	Chen, Y., Lebrun, J.-J., and Vale, W. (1996). Regulation of transforming growth factor $\beta$ - and activin-induced transcription by mammalian Mad proteins. Proc. Natl. Acad. Sci. USA 93, 12992-12997.
	7.	Dennler, S., Itoh, S., Vivien, D., ten Dijke, P., Huet, S., and Gauthier, J.-M. (1998). Direct binding of Smad3 and Smad4 to critical TGF $\beta$ -inducible elements in the promoter of human plasminogen activator inhibitor-type 1 gene. EMBO J. 17, 3091-3100.
	8.	Dyson, S. and Gurdon, J.B. (1998). The Interpretation of Position in a Morphogen Gradient as Revealed by Occupancy of Activin Receptors. Cell 93, 557-568.
	9.	E. Labbé et al., (1998), Molecular Cell, v. 2, pp. 109-120.
	10.	EMBL Sequence Database Accession No. EMHUM2.AB002303, KIAA0305
	11.	EMBL Sequence Database, Accession No. EMHUM1:AF130419
	12.	Faux, M. and Scott, J.D. (1996). Molecular glue: kinase anchoring and scaffold proteins. Cell 85, 9-12.
	13.	Gaullier et al., (1998), Nature, 394, 432-433.
	14.	Heldin, C.-H., Miyazono, K., and ten Dijke, P. (1997). TGF- $\beta$ signaling from cell membrane to nucleus through SMAD proteins. Nature. 390, 465-471.
	15.	Henis, Y.I., Moustakas, A., Lin, H.Y., and Lodish, H.F. (1994). The type II and III transforming growth factor- $\beta$ receptors form homo-oligomers. J. Cell Biol. 126, 139-154.
	16.	Hoodless, P.A., Haerry, T., Abdollah, S., Stapleton, M., O'Connor, M.B., Attisano, L., and Wrana, J.L. (1996). MADR1, a MAD-related protein that functions in BMP2 signaling pathways. Cell 85, 489-500.
	17.	Kim, J., Johnson, K., Chen, H.J., Carroll, S., and Laughon, A. (1997). <i>Drosophila</i> Mad binds to DNA and directly mediates activation of <i>vestigial</i> by decapentaplegic. Nature 388, 304-308.
	18.	Kretzschmar, M. and Massagué, J. (1998). SMADs: mediators and regulators of TGF- $\beta$ signaling. Current Opinion in Genetics & Development 8, 103-111.
123	19.	Kretzschmar, M., Liu, F., Hata, A., Doody, J., and Massagué, J. (1997). The TGF- $\beta$ family mediator Smad1 is phosphorylated directly and activated functionally by the BMP receptor kinase. Genes Dev. 11, 984-995.

EXAMINER

DATE CONSIDERED

\*EXAMINER

Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.


FORM PTO-1449 U.S. Department of Commerce  
Patent and Trademark Office

Attorney Docket Number  
3477.91

Serial No.  
09/744,167

LIST OF DOCUMENTS CITED BY APPLICANT

(Use several sheets if necessary)



RECEIVED

SEP 30 2002

Applicants:

Jeffrey L. Wrana

TECH CENTER 1600/2900

Filing Date  
January 19, 2001

Group

20.	Labbe, E., Silvestri, C., Hoodless, P.A., Wrana, J.L., and Attisano, L. (1998). Smad2 and Smad3 positively and negatively regulate TGF $\beta$ -dependent transcription through the forkhead DNA binding protein, FAST2. <i>Molecular Cell</i> in press.
21.	Lagna, G., Hata, A., Hemmati-Brivanlou, A., and Massagué, J. (1996). Partnership between DPC4 and SMAD proteins in TGF- $\beta$ signaling pathways. <i>Nature</i> 383, 832-836.
22.	Liu, X., Sun, Y., Constantinescu, S.N., Karam, E., Weinberg, R.A., and Lodish, H.F. (1997b). Transforming growth factor $\beta$ -induced phosphorylation of Smad3 is required for growth inhibition and transcriptional induction in epithelial cells. <i>Proc. Natl. Acad. Sci. USA</i> 94, 10669-10764.
23.	M. Kretschmar et al., (1998), <i>Current Opinion in Genetics &amp; Development</i> , v. 8, pp. 103-111.
24.	Macías-Silva, M., Abdollah, S., Hoodless, P.A., Pirone, R., Attisano, L., and Wrana, J.L. (1996). MADR2 is a substrate of the TGF $\beta$ receptor and its phosphorylation is required for nuclear accumulation and signaling. <i>Cell</i> 87, 1215-1224.
25.	Nagase et al., (1997), <i>DNA Research</i> , v. 4, pp. 141-150.
26.	Nakao, A., Imamura, T., Souchelnytskyi, S., Kawabata, M., Ishisaki, A., Oeda, E., Tamaki, K., Hanai, J.-i., Heldin, C.-H., Miyazono, K., and ten Dijke, P. (1997a). TGF- $\beta$ receptor-mediated signaling through Smad2, Smad3 and Smad4. <i>EMBO J.</i> 16, 5353-5362.
27.	Nakayama, T., Snyder, M.A., Grewal, S.S., Tsuneizumi, K., Tabata, T., and Christian, J.L. (1998). <i>Xenopus</i> Smad8 acts downstream of BMP-4 to modulate its activity during vertebrate embryonic patterning. <i>Development</i> 125, 857-867
28.	Nishimura, R., Kato, Y., Chen, D., Harris, S.E., Mundy, G.R., and Yoneda, T. (1998). Smad5 and DPC4 are key molecules in mediating BMP-2-induced osteoblastic differentiation of the pluripotent mesenchymal precursor cell line C2C12. <i>J. Biol. Chem.</i> 273, 1872-1879.
29.	Patki et al., (1998), <i>Nature</i> , 394, 433-434.
30.	Patterson, G., Kowek, A., Wong, A., Liu, Y., and Ruvkun, G. (1997). The DAF-3 Smad protein antagonizes TGF- $\beta$ -related receptor signaling in the <i>C. elegans</i> dauer pathway. <i>Genes and Dev.</i>
31.	Pawson, T. and Scott, J.D. (1997). Signaling through scaffold, anchoring and adaptor proteins. <i>Science</i> 278, 2075-2080.
32.	Reddy, A.H. (1998), <i>Nature Biotechnology</i> , v. 16, pp. 247-252

EXAMINER  
\*EXAMINER

DATE CONSIDERED

Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

[illegible]

EXAMINER  
\*EXAMINER

DATE CONSIDERED

Initial if reference considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.